Long lasting breeding performance differences between wild-born and released females in a reinforced North African Houbara bustard (*Chlamydotis undulata undulata*) population: a matter of release strategy

Leo Bacon, Alexandre Robert, Yves Hingrat

► To cite this version:

Leo Bacon, Alexandre Robert, Yves Hingrat. Long lasting breeding performance differences between wild-born and released females in a reinforced North African Houbara bustard (*Chlamydotis undulata undulata*) population: a matter of release strategy. *Biodiversity and Conservation*, 2019, 28 (3), pp.553-570. 10.1007/s10531-018-1651-6 . mnhn-02553975

HAL Id: mnhn-02553975

<https://hal-mnhn.archives-ouvertes.fr/mnhn-02553975>

Submitted on 24 Apr 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Long lasting breeding performance differences between wild-born and released females in a reinforced North African Houbara bustard (*Chlamydotis undulata undulata*) population: a matter of release strategy.

Léo Bacon^{a, b, *, **}, Alexandre Robert^b, Yves Hingrat^{a, c}

^aEmirates Center for Wildlife Propagation, PO Box 47, 33250 Missour, Morocco.

^bCESCO, UMR7204 MNHN-CNRS-Sorbonne Université, CP135, 43 Rue Buffon, 75005 Paris, France.

^cReneco International Wildlife Consultants LLC., Po Box 61741, Abu Dhabi, U.A.E.

*Corresponding author (bacon.leo@gmail.com)

**Orcid ID: 0000-0003-1200-9902

Abstract

The success of translocation programmes is reflected by the ability of translocated individuals to survive and reproduce in their new environment. However, it has previously been reported that translocated individuals have lower demographic performance than their wild-born conspecifics, due to management and individual factors (such as release conditions or age). Here, we study six breeding parameters in free-ranging females of the North African Houbara bustard (*Chlamydotis undulata undulata*) and compare these parameters between captive-bred released (n=204) and wild-born (n=101) birds, considering the age of individuals and the period of release (autumn *versus* spring). Our results indicate that (1) captive-bred released females successfully breed in the wild; (2) for three out of the six breeding parameters studied, released females show lower performances than wild-born females; but, (3) Although we observed consistently reduced breeding performances in one year old females relative to older females,

we did not uncover any interaction between age and the origin of females, suggesting that the impairment of breeding parameters in released females is long lasting; and, (4) interestingly, this impairment of breeding parameters depends on the period of release, with lower breeding performances for spring releases compared to autumn releases. Overall, our study highlights the capacity of captive-bred females to reproduce in the wild, contributing to the dynamics of the population beyond their individual history. Our results also uncover complex variations of breeding parameters in translocated birds, but suggest that these differences can be minimized through an appropriate translocation strategy.

Key-words: Captive-breeding, post-release effect, reinforcement, reproduction, translocation

Introduction

The success of conservation translocation programmes is commonly defined as the capacity of translocated populations to persist without further intervention (Ewen et al. 2012, IUCN 2013). Assessment of success thus requires an understanding of the long-term dynamics and viability of translocated populations (Robert et al. 2015a), which first and foremost requires a long-term demographic assessment (Sutherland et al. 2010). However, such assessments are rarely achieved, mainly due to a lack of resources to support appropriate monitoring and difficulties associated with data analysis (Fischer & Lindenmayer 2000, Sutherland et al. 2010). Translocated populations are complex, heterogeneous and unbalanced systems (Robert et al. 2007), and the assessment of demographic performances is required to encompass this complexity. In particular, it is necessary to assess potential demographic variations among translocated and wild-born organisms.

Previous knowledge indicates that the demography of translocated populations is likely to be affected by a variety of genetic and non-genetic issues. Genetic issues include outbreeding depression (Huff et al. 2011) and ill adaptation (Montalvo & Ellstrand 2001) as well as

inbreeding and drift loads (Robert 2009). In cases of translocations based on captive breeding, adaptation to captivity (Frankham 2008) might be an important additional issue. In animals, non-genetic issues are associated with captive-rearing and release strategies, leading to various factors potentially affecting demographic rates in the wild. These include individual condition, such as health status, physiology and behaviour (Champagnon et al. 2012, Dickens et al. 2010, Hardouin et al. 2014, Tavecchia et al. 2009), proximity of the release site with other populations (Mihoub et al. 2011), period of release (Hardouin et al. 2014), age at release (Sarrazin et al. 1996) or the size and composition of the animal group (Hardouin et al. 2014, 2015a). Additionally, the potential impairment of demographic performances of translocated individuals might be explained by the interaction of their phenotype with their new environment and release conditions.

Post-release effects have been defined as impairments of vital rates caused by translocation conditions (Ewen et al. 2012). Such effects can be estimated either by assessing changes in vital rates over time in a release cohort (e.g., Armstrong & Ewen 2001, Tavecchia et al. 2009) or by comparing vital rates of translocated and resident animals over the same time period (e.g., Brown et al. 2006). A number of studies in translocated vertebrates have documented these post-release effects, mostly in terms of survival probabilities (Armstrong et al. 2017, Bertolero & Oro 2009, Hardouin et al. 2014, Sarrazin et al. 1994) but also in reproduction rates (Bertolero & Oro 2009, Converse et al. 2013, Sarrazin et al. 1996, Tavecchia et al. 2009) and even in dispersal behaviour (Le Gouar et al. 2008, Mihoub et al. 2011). In the particular cases of translocated individuals originating from captive-breeding, any impairments of vital rates in translocated individuals can result either from translocation conditions (i.e., post-release effects *per se*), captivity conditions (including genetic and non-genetic issues), or the interaction of translocation and captivity conditions.

However, disentangling the various sources of demographic variation in translocated populations (and in particular those related to the translocation protocol from other sources of variation) is challenging because (1) the comparison between translocated and wild-born animals requires the monitoring of wild-born individuals; a difficult task often hindered by several methodological and ethical issues, and (2) changes of vital rates over time in translocated individuals can be confounded by other longitudinal sources of demographic heterogeneity, such as age effects (Bacon et al. 2017a), or by inter-individual heterogeneity (e.g., apparent improvement of survival due to the selection of the best survivors).

The North African Houbara bustard (*Chlamydotis undulata undulata*, hereafter Houbara – Fig. 1) is a medium-sized bird historically distributed from Northern Mauritania to Egypt. It is threatened by over-hunting, poaching and habitat degradation, which have led to the decline of the species during the second half of the twentieth century (Goriup 1997). This decline led to the establishment of a reinforcement programme (the Emirates Center for Wildlife Propagation, ECWP, see Material and Methods) which employs captive breeding and regular releases of captive-bred individuals, combined with ecological research and hunting management, in order to increase the population size of the threatened Houbara throughout its range and to maintain viable populations in adequacy with traditional Arab falconry (Lacroix et al. 2003).

Importantly, captive-bred Houbara are released as juveniles (within their first year of life) either before or during the breeding season (in autumn and in spring, respectively). Previous researches on translocated Houbara indicate that (1) captive-bred juveniles exhibit reduction in their short-term survival (the magnitude of which varies depending on the release season and meteorological conditions), compared to long-term survival (Hardouin et al. 2014), and (2) one-year-old released females have reduced breeding performance relative to older released females (Bacon et al. 2017a). These results suggest that translocated individuals may experience changes of vital rates over time. However, in these previous assessments of demographic rates,

potential changes caused by translocation conditions, captivity conditions and their interactions may have been confounded with age effects. Thus, a rigorous assessment of these potential captivity and translocation effects requires formal comparison between captive-bred released and wild-born individuals of different ages. Hereafter, demographic effects occurring only in the first year following release (i.e., in one-year-old individuals) are referred to as short-term differences and effects occurring independently of the time since release (i.e., in individuals of any age) are referred to as long lasting differences.

Here, we investigate potential differences in breeding performances between wild-born and released captive-bred females, while controlling for age, period of release, and temporal and individual effects. The Houbara is a ground-nesting, gyneparental incubating species. In Eastern Morocco, females breed from mid-February to mid-June. They lay generally two to three eggs, at intervals of 2.5 days, which they incubate for an average of 23 days (Gaucher 1995). In case of nest failure, females generally initiate a replacement clutch. Houbara chicks are nidifugous but still rely on their mother for food during the first 10 days after hatching (Saint Jalme & van Heezik 1996) and fledge at a mean age of 60 days (Hardouin et al. 2012). Maximum longevity observed in the wild was at least 13 years old for a wild-born female and 10 years old for captive-bred released females (Bacon et al. 2017a). The diet of the Houbara bustard is generalist and opportunistic and present strong seasonal variation in animal (e.g. Coleoptera, Hymenoptera) and vegetal (e.g. Asteraceae, Brassicaceae, Chenopodiaceae – Bourass et al. 2012) proportion.

We used data collected from a longitudinal nest survey conducted between 2002 and 2016 in the reinforced Houbara population in Eastern Morocco.

We focused on six breeding parameters: nesting effort (number of nesting attempts per breeding season per female), nest initiation date, clutch size, egg volume, daily nest survival and brood

survival. We hypothesized that potential short-term reduction of performance in released Houbara are related to their potentially low condition and limited experience with their new environment (Bertolero & Oro 2007, Bertolero et al. 2009). Thus, potential differences between released and wild-born individuals should decrease with increasing age, reflecting both individual (i.e., gain in experience within an individual, Mauck et al. 2012) and population-scale processes (i.e., death of lower-quality individuals, assuming that survival and breeding performance may positively covary, Robert et al. 2015b). We thus hypothesize that the difference in breeding performances of wild-born and captive-bred released females will decrease with their age (which is confounded with the time since release for released females). In other words, we expect to find an interaction between the origin of females (released vs. wild-born) and their age.

Material and methods

Study area

The study was conducted in Eastern Morocco, in the ECWP intervention area (Fig. 2). The study area encompasses approximately 50 000 km² and is characterized by an arid climate marked by irregular rainfalls (less than 200 mm/year). Mean temperatures vary from 6.80°C in winter (December to February) to 26.64°C in summer (June to August). The habitat in the study area is characterized by sparse, shrubby vegetation. Vegetation cover in the plains mainly comprises Chenopodiaceae such as *Salsola* spp., *Hamada* spp. and Compositae such as *Artemisia herba-alba*. In clay and silt-rich areas created by seasonal runoff waters, *Salsola* sp. is often associated with *Atriplex* spp. Drainage courses and wadis are characterized by *Zizyphus lotus* (Rhamnaceae) and *Retama* sp. (Leguminosae). On the high plateaux and slopes, from the extreme east to the Oran region of Algeria, the vegetation is dominated by *Stipa tenacissima* (Gramineae). The core of the study area is used for nomadic pastoral activities, herding and

144 grazing. Mixed herds are generally composed of 100–300 heads of sheep and goats. Nomads
145 settle camps within the study area, whereas some shepherds come from surrounding villages
146 (Le Cuziat et al. 2005).

147 *Emirates Center for Wildlife Propagation and reinforcement programme*

148 The EWCP was established in 1995 to mitigate the severe decline in Houbara populations
149 occurring in the second half of the 20th century. Two ECWP breeding stations were built for
150 captive breeding (in Missouri in 1995 and in Enjil in 2005, see online appendix A). Detailed
151 descriptions of the captive breeding programme can be found in Lesobre et al. (2010) and
152 Chargé et al. (2010, 2014). From 1996 to 2016, 108 486 birds were released in North Africa
153 (94 374 in the study area). Locally, these intensive releases may induce density dependent
154 process leading to deleterious biotic interactions (intraspecific competition, disturbance,
155 predation) and negatively impact life history parameters (Azar et al. 2016, Bacon et al. 2017b).

156 Within the study area, the overall number of birds released progressively increased over the
157 years (from 28 in 1996 to 9 084 in 2016, see online appendix B), with an increase in the number
158 of release sites and a decrease in the group size per release (Hardouin et al. 2014). The release
159 group size varied between sites (from four to 498 individuals, median = 18 individuals).
160 Houbara were released at an average of 6 ± 3 (standard deviation) months of age in autumn
161 (August to December), and an average of 9 ± 1 months of age in spring (February to May).
162 Autumn releases were managed to avoid direct mortality and disturbance due to hunting. In
163 addition, since 2003, some birds were released in summer (June) at an average of 3 ± 0.6 months
164 of age, but they represent only 8% of total releases in the study area. As such, in this study, we
165 only focused on captive-bred females released in spring and autumn.

166 Hunting was banned between 2000 and spring 2005, and subsequently restricted to the
167 autumn/winter period (October–January) confined within an area comprising 60% of the

intervention area. Since 2014, it is estimated that on average 2000 birds are harvested every year, with captive-bred individuals representing 85% of the hunting bags (ECWP unpublished data).

Monitoring of captive-bred released females

Prior to 2006, released Houbara were tagged on the left tarsus with an aluminium ring displaying a unique ID number. Beginning in 2006, all released individuals were subcutaneously tagged with a unique radio-frequency identification (RFID) microchip (TROVAN LID100 implantable transponder, DorsetID; see Hardouin et al. 2015b for details). In Eastern Morocco, 3 014 released birds (49% females) were equipped with transmitters (1 937 with VHF transmitters, 933 with satellite transmitters and 144 with GSM transmitters). The VHF transmitters used in this study were battery-powered necklace units with mortality signals (11 g model RI-2B-M, 20 g model RI-2D-M, Holohil System Ltd., Carp, Ontario, Canada) and solar-powered backpack (19 g model, Merlin Systems Inc., Boise, Idaho, USA). During the breeding season, females fitted with VHF transmitters were searched for at least once per week. The satellite and GSM transmitters used in this study were backpack solar-powered transmitters: PTT-100 30 or 45 g Solar GPS PTT and 30 g solar GSM (Microwave Telemetry, Inc., Columbia, Maryland, USA). To identify breeding events of individuals equipped with satellite transmitters, individuals' movements were remotely monitored. When stationary locations were recorded, the last location was checked in the field to confirm whether the individual was dead or nesting.

Trapping and identification of wild-born nesting females

Trapping of wild-born females was performed during egg-collection campaigns in the breeding season of the periods 1997-1998, 2001-2009 and 2015-2016 (see details in online appendix C). Egg collection campaigns were aimed at building the founder population of the ECWP captive-

breeding program (Lesobre et al. 2010, Chargé et al. 2014). Females were caught using nylon snares around their nest or around their recently hatched chicks. In total, 143 wild-born females were caught while nesting and were equipped with a transmitter or transmitter replacement. In addition, 10 wild-born females captured and equipped with VHF/PTT transmitters as juveniles were found nesting in later years (see Hardouin et al. 2012 for more details of the capture and monitoring procedures).

Data collection

Breeding parameters

Six breeding parameters were assessed from breeding survey data collected from 2002-2016: the nesting effort (number of nesting attempts per breeding season per female), the clutch size, the egg volume, the nest initiation date, the daily nest survival rate and the daily brood survival rate. The number of years for which survey data was available varied slightly between breeding parameters, this is detailed in online appendix D.

- Nest survey

Nests were located from a collaborative survey with shepherds and from individually tracked females. Because of the bias inherent to the trapping and egg-collection campaigns (all wild-born non-equipped nesting females had their eggs collected if not already hatched and were caught) and to avoid confounding factors between transmitter presence and female origin (wild-born and captive-bred), we only selected those nests for which the nesting female was equipped with a transmitter at nest initiation. In total, 703 nests from 305 females were selected for this study. For wild-born females, 300 nests were from 91 wild-born females first captured on their nest, and 21 were from 10 females captured and marked at the juvenile stage. For captive-bred females (n nests = 382), we selected nests from cohorts of females that were released in their

first year of life from August to November (in autumn, before the breeding season, n nests = 122, n females = 74) and from February to May (in spring, during the breeding season, n nests = 260, n females = 130). Nests were visited on average every 6 ± 3.60 days. In addition, since 2013, camera traps were deployed at some of the monitored nests (n=45), providing a continuous survey effort and more accurate assessments of nest fate. Nests equipped with camera traps were visited on average every 8 ± 5.07 days. Preliminary unpublished results did not highlight any deleterious effects of the camera trap monitoring on nest survival in our study area (Bacon 2017a). The use of camera traps gave great details on the guild of species predating on Houbara nests. It is composed of the red fox (*Vulpes vulpes*), the golden jackal (*Canis aureus*), stray dogs (*Canis lupus familiaris*), desert hedgehog (*Paraechinus aetiopicus*), common raven (*Corvus corax*) and brown-necked raven (*Corvus ruficollis*). Nest failure was recorded for a visit interval if the clutch disappeared without signs of hatching (Mabee 1997) or was abandoned. We considered a nest to have survived if at least one egg hatched (Mayfield 1975).

The clutch size was defined as the maximum number of eggs observed in the nest. Since 2003, the length, width (to the nearest hundredth of a centimetre) and mass (to the nearest tenth of a gram) of each egg were measured. The width and length of eggs were then used to calculate the egg volume (Vol) in cm^3 (see details in Bacon et al. 2017a). The nest initiation date was estimated from the calculation of the eggs' incubation stage (in days) at the time of measurement (based on the egg weight loss equation from Hoyt 1979, see details in Bacon et al. 2017a).

The nesting effort was calculated as the number of nests initiated per breeding season per female, including females initiating no nest. We selected females that were continuously monitored during the breeding season. Even if all females were fitted with monitoring devices, technical limitations could cause gaps in their monitoring, which could cause the non-detection

of nesting attempts. To increase the detectability of nesting attempts, we only selected females that were fitted with VHF and monitored from February to the end of June, with at least one individual sighted every two weeks (resulting in 173 monitoring histories from 141 females from 2002 to 2014).

Trapping of nesting females causes systematic nest abandonment; thus, female trapping and egg collection have a positive effect on overall nesting effort, as they promote replacement clutches. However, replacement clutches are initiated later in the breeding season, affecting the distribution of nest initiation dates. Therefore, for those two parameters, we excluded females from the years where their eggs were collected (see online appendix D).

- Brood survey

Assessing breeding success after hatching and before fledging in species with nidifugous chicks is difficult. It requires the marking and close monitoring of all individuals; a delicate operation that may affect chick survival at this very sensitive stage of life (Barron et al. 2010). As such, we chose to not mark chicks and instead assessed the fate of broods through the monitoring of brooding females. Therefore, between 2002 and 2014, nesting females fitted with VHF transmitters that successfully hatched at least one egg were then searched for at least once per week. Brood failure was considered for a female at a given monitoring interval if the entire brood disappeared before 60 days of age (Hardouin et al. 2012). In total, we monitored 201 broods from 133 females. A few females (n=12) were captured while rearing chicks, at the nest or after leaving the nest, either to take measurements or to replace the transmitter.

Statistical analyses

Nest and brood survival analyses

We used the logistic exposure approach of Shaffer (2004) to estimate the daily nest survival rate (defined as the probability of a living nest to not fail in one day). Egg collections and female trapping cause the end of the nest history and produce an unknown “natural” final fate. Theoretically, in occasional cases where they represent a subset of the sample of nests monitored, nests with an unknown final fate are right truncated at the last date they are known to be active for daily nest survival analysis. In our study, the number of clutches collected was relatively important and biased towards wild-born females. Therefore, nests with an unknown final fate were not representative of the sample of the nests monitored and could lead to an overestimation of the daily nest survival of wild-born females relatively to released females. Thus, we excluded all nests with an unknown final fate from the analysis (following Manolis et al. 2000 and personal communication from Stephen Dinsmore). Finally, as records of periodic nest visits were only standardized beginning in 2003, we discarded data collected prior to 2003.

The same statistical approach was used to study brood survival, with the exception that the exposure period starts at hatching and ends 60 days later for successful broods (mean age at fledging, Hardouin et al. 2012, but see details in Bacon et al. 2017a).

Clutch size, egg volume, nest initiation date and nesting effort

The nesting effort was analysed using a generalized linear mixed effects model with a Poisson distribution and a log link function. The clutch size was analysed based on ordinal regression using cumulative linked mixed-effects models (CLMMs). Egg volumes and nest initiation dates were analysed using linear mixed-effects models (LMMs) fitted by maximum likelihood with a Gaussian distribution and an identity link function.

Origin and age factor

We grouped nests into categories based on female origin (wild-born, captive-bred released in autumn and captive-bred released in spring). In addition, we assigned to each nest the age of the nesting female: one-year-old or at least two years old (see online appendix E). For wild-born females, only those captured as juveniles could be aged precisely. A previous study focussing only on released females indicated that one-year-old females had lower breeding performance than older individuals (Bacon et al. 2017a). In addition, the percentage of one-year-old released females initiating at least one nest is low (8%, ECWP unpublished data). Based on this information, we assumed that all wild-born females with unknown age were at least two years old at their first identification on a nest.

Adjustment variables

In monitored animals, events such as trapping and marking are stressful and may have some effect on subsequent behavioural and demographic patterns (e.g. unusual movements, increased mortality, cessation of breeding) once released (Casas et al. 2015, Ponjoan et al. 2008). For nesting female Houbara, trapping causes nest abandonment (direct effect) but may also have a carry-over effect on subsequent breeding performance. For females trapped on live chicks, trapping may have a direct effect on brood survival by affecting either the female or the chicks. Therefore, we considered an intra-seasonal carry-over effect of trapping for the following parameters: clutch size, egg volume, daily nest survival and daily brood survival rate, when measured on the replacement clutch consecutive to the trapping of the female (carry-over trapping). In addition, specifically for the daily brood survival rate, we considered a direct effect of trapping (direct trapping) on the brood when trapping of the females occurred while rearing chicks. Carry-over trapping and direct trapping effects were implemented as binary adjustment variables. To account for the temporal variation of breeding performance values within the breeding season, we included a date covariate (in Julian days, 1st of January = 1, implemented as the nest initiation date for egg volume and clutch size, as the median date between two nest

visits for the daily nest survival rate, and as the hatching date of the nest for the daily brood survival rate, Grant & Shaffer 2012). Finally, we considered temporal variation of the daily survival rate of the nest and brood by including, respectively, a binary factor describing whether the nest was in laying or incubation stage and a continuous covariate indicating the age of the brood since hatching (Grant & Shaffer 2012).

Model selection

For each breeding parameter analysis, covariates were rescaled, centralized and standardized by two times the standard deviation (Gelman 2008), and factors with two levels were transformed into binary dummy variables (0-1).

We examined the relationship between the breeding parameters and their corresponding variables by developing sets of candidate models that included additive terms of explanatory variables, quadratic terms of continuous temporal covariates (date² and brood age²) and the interaction between the age of the nesting female and its origin. To adjust for potential effects of trapping, we kept the trapping carry-over effect and the direct trapping effect in all candidate models of the concerned breeding parameters. Additionally, current year, female identity and nest identity nested under female identity (only for egg volume analyses) were incorporated as random intercept effects in all models. Random intercept effects were included to avoid pseudo-replication and to account for potential heterogeneity among years, females and nests (Zuur et al. 2009).

Thus, the starting (most complex) statistical models were of the following form:

$$Y = \text{Female Age} + \text{Female Origin} + \text{Female Age} \times \text{Female Origin} + \text{Adjustment variables} \\ + \text{random intercepts}$$

where Y was the dependent variable (the breeding parameter). Details of specific adjustment variables (mentioned in the section *Adjustment variables*) implemented in the starting model of each breeding parameter can be found in Appendix F.

Starting from this generic structure, models were simplified to develop all possible combinations of explanatory variables (including additive, polynomial terms and the interaction, see all model combinations in Appendix G). This procedure was applied to our six breeding dependent variables. For each breeding dependent variable, we ranked models using the Akaike Information Criterion corrected for small sample size (AICc). The best model ($\Delta\text{AICc} = 0$; Burnham & Anderson 2002) was selected to estimate β coefficients and associated 95% confidence intervals for fixed-effect covariates. All analyses were conducted using R 3.3.3 (R Development Core Team, 2017) and the packages lme4 1.1-12 (Bates et al. 2015), Ordinal 2015.6-28 (Christensen 2015), MuMIn 1.13.4 (Barton 2015), and nest survival (Herzog 2009). Normality and homoscedasticity of the random residuals were checked graphically from the starting models, and no overdispersion (Zuur et al. 2009) was observed for the Poisson distribution model.

Results

On average, females were monitored over 2.97 ± 1.50 years (3.58 ± 1.76 years for wild-born females and 2.70 ± 1.25 years for released females) and 86% of females were monitored over more than 1 year (88% for wild-born females and 85% for released females).

We analysed the effects of dependent covariates on our six breeding parameters based on a common statistical approach. To keep the presentation of the results clear and straightforward, we directly present the coefficients from the best selected models ($\Delta\text{AICc} = 0$) in Table 1. AICc based model ranking selections, which compiled six tables, can be found in online appendix G.

According to the best models (Table 1), clutch size and egg volume declined along the breeding season. Daily nest survival was higher during the nest initiation stage than during the incubation stage, while daily brood survival increased with the age of the brood. For all breeding parameters concerned (clutch size, egg volume, daily nest survival and daily brood survival), the carry-over trapping effect had 95% confidence intervals overlapping 0, whereas direct trapping of females while rearing chicks had a very strong negative impact on the daily survival of the brood (Table 1). For all breeding parameters (with the exception of the daily nest survival rate), the Age effect was selected in all best models, indicating that one-year-old females had poorer breeding performances than older ones (Table 1).

The origin factor was selected in the best model sets for nesting effort, egg volume and daily brood survival rate. In Table 1, for the female origin factor, the results show the differences with respect to a reference level at the intercept. We defined wild-born females as the reference level at the intercept in Table 1, but all levels of the female origin factor were set as the reference level at the intercept in order to assess potential significant differences. Captive-bred females initiated fewer nests and produced smaller eggs than wild-born females (Figs 3A, B). However, confidence intervals of these effects did not overlap 0 only for captive-bred females released in spring relative to wild-born females (Table 1). Captive-bred females released in autumn had intermediate values of nesting effort and egg volume, but with confidence intervals difference to the two other groups overlapping 0 (Table 1). Finally, captive-bred females released in spring had lower daily brood survival than wild-born females and captive-bred females released in autumn (Fig. 3C, Table 1), and the confidence intervals of the difference between captive-bred females released in spring and wild-born females was close to negative (Table 1).

We were particularly interested in the interaction term of the age of nesting females and their origin, which quantifies whether any effect of origin depends on age (i.e., short-term) or not (i.e., long lasting). However, none of the best models (Table 1) included this interaction,

providing no support for a more marked effect of the origin in one-year-old females as compared to older females.

Discussion

The assessment of the breeding performances of released individuals is of major importance in translocation programmes (Bertolero et al. 2009). In this study, we investigated the differences of breeding parameters between wild and captive-bred released Houbara females and revealed how release strategies may influence the magnitude of this differences. This is a prerequisite to adapting and improving translocation strategies and thus to increase the chance of translocation success (Armstrong & Seddon 2008).

Most breeding parameters (except nest survival) consistently showed age-specific variation, suggesting that one-year-old females have reduced performances compared to older ones, in agreement with a previous study (Bacon et al. 2017a). Individual heterogeneity was taken into account in our statistical framework (see Nussey et al. 2006, van de Pol & Verhulst 2006). This suggests that the observed effect of age is best explained by longitudinal improvement of individuals (increased breeding experience or an increase in breeding investment, Curio 1983), than by population level processes (e.g., selection of the best individuals over time). The lack of age effects on the daily nest survival may be explained by the weight of external factors, such as environmental conditions (Morales et al. 2002) or predation, which is a prominent cause of nest failure in birds (Martin 1995), including Houbara (Bacon et al. 2017a, Koshkin et al. 2016), potentially overriding any age-dependent effects.

Regardless of age effects, our results indicate that, for some of the breeding parameters studied, wild-born individuals performed slightly better than individuals translocated in autumn and better than individuals translocated in spring. More precisely, wild-born females initiated more nests with larger eggs when compared to captive-bred females released in spring. In addition,

the daily brood survival of captive-bred females released in spring was lower than for wild-born females. These differences have a marked size effect. It corresponds to a decline of ~47% in nesting effort and a decline of ~40% in brood survival (from hatching to fledging) when considering adult females released in spring relative to wild-born adult females (Figs 3A, C).

Conversely to our expectations, we did not observe any interaction between the age of the female and its origin. This result is consistent with the **long lasting** (i.e., permanent) negative post release effect on breeding parameters observed in other translocated species (see, e.g., Sarrazin et al. 1996). It is also consistent with previous findings in Houbara that released captive-bred females exhibit age-dependent patterns of variation in breeding performance, similar to those observed in many vertebrate populations in the wild (i.e., a bell-shaped trend divided into youth, adulthood and senescence; see discussion in Bacon et al. 2017a).

As differences between wild-born and captive-bred individuals were **long lasting**, we cannot exclude the possibility that genetic causes related to captive-breeding issues partly explain the observed differences. However, strong and restrictive guidelines are conducted in the ECWP captive breeding programme (Chargé et al. 2014). These guidelines aim to minimize the loss of genetic variation and to minimize the adaptation to captivity in the captive population in order to maximize the fitness of individuals released into the wild, while reducing the genetic consequences of the reinforcement programme on wild populations (Chargé et al. 2014). Nevertheless, the relaxation of sexual and natural selection and adaptation to captivity may partly explain the lower breeding performances of captive-bred individuals in the wild (Araki et al. 2007, Frankham 2008). Beside, owing to the substantial translocation effort (see online appendix B), a substantial proportion of wild-born individuals may be offspring from translocated birds (see general discussion in Barbanera et al. 2010), which may explain why the breeding performances of released and wild-born females are relatively similar. Therefore, it is likely that most differences between the two groups are related to non-genetic issues.

At the phenotypic level, previous research into captive breeding/rearing in birds has suggested that demographic differences between released and wild-born individuals can be explained by the lack of exposure of chicks hand-reared in captivity to adult behaviour (Roche et al. 2008), permanently decreasing their fitness in the wild (Buner & Schaub 2008, Buner et al. 2011, Pérez et al. 2015, Sánchez-Garcia et al. 2011). Although wild-born and released captive-bred Houbara females have a similar diet (Bourass & Hingrat 2015) and exploit equivalent ecological niches (Monnet et al. 2015), released females may be less efficient at finding and exploiting resources (Champagnon et al. 2012). This could explain the lower investment of captive-bred females in reproduction (e.g., egg production), which may subsequently affect the survival of the chicks. In species with precocial chicks (including nidifugous chicks), smaller eggs generally contain less yolk, giving rise to chicks that are in lower condition with lower nutrient reserves at hatching and that are more prone to early mortality (Amundsen & Stokland 1990, Blomqvist et al. 1997, Galbraith 1988, Williams 1994). The differences observed between wild-born and translocated females must be interpreted while taking into account that all individuals were fitted with transmitters. Markings devices can have significant impacts on some demographic, physiological and behavioural parameters (Barron et al. 2010, see discussion in Le Maho et al. 2014). However, because wild-born and released females are unlikely to exhibit differential responses to transmitter fitting, our main conclusions are unlikely to be affected by transmitter effects.

Our results indicate that the difference observed between wild-born and released females varies according to the period of release (autumn versus spring). Captive-bred females released in autumn had consistently higher nesting effort, egg volume and daily brood survival values than females released in spring (Table 1). Assuming that survival and reproductive parameters positively covary (e.g., "fixed heterogeneity", Cam et al. 2016), such a pattern may be related to the death of the less performant females between their release and the breeding season among

females released in autumn. This hypothesis is supported by previous findings showing that harsh environmental conditions in autumn and winter lower short-term survival rates compared to spring releases (Hardouin et al. 2014), representing a stronger environmental filter that primarily affects weak individuals.

It is important to note that not all breeding parameters differed between wild-born and captive-bred females. The origin of females was not selected in the best models for clutch size, nest initiation date and daily nest survival analysis. Similar heterogeneous responses of multiple breeding parameters to female origin were observed by Hill & Roberston (1988) for a reinforced ring-necked pheasant (*Phasianus colchicus*) population, where wild-born and captive-bred released females had similar nesting effort and nesting success, yet chick survival to fledging and total productivity were higher for wild-born females.

Management implications

We observed consistent, permanent impairments in some breeding parameters of captive-bred released females relative to wild-born females. The magnitude of these differences highlights the importance of assessing demographic impacts of captivity and post release effects in translocated populations. In our study, the observed differences between wild-born and released females appears to be dependent on the translocation strategy (period of release), which implies that it could be buffered by adapting translocation strategies (favouring autumn releases). Nonetheless, the current strategy (spring and autumn releases) seems suitable regarding the multiple objectives of the reinforcement programme, i.e., reinforcement of the wild population and supplementation for hunting (falconry). For autumn releases, environmental stochasticity may select individuals in good condition (Hardouin et al. 2014) that will later breed with relatively better parameters (as suggested by the present results). On the other hand, although outcomes from spring releases will be less certain in terms of breeding performances, higher

survival rates in this season will help to maintain the population size at a level that can sustain hunting pressure (Hardouin et al. 2015a) and population growth rate. Finally, although differences were observed between wild-born and captive-bred females, our results confirm the capacity of captive-bred females to reproduce in the wild and to contribute to the dynamics of the population beyond their individual history. This partially supports the success of the reinforcement programme (Bertolero & Oro 2009) and provides a strong basis for more integrative and quantitative assessment of translocation success (Robert et al. 2015a).

Acknowledgements

This study was funded by the Emirates Center for Wildlife Propagation (ECWP), a project of the International Fund for Houbara Conservation (IFHC). We are grateful to H.H. Sheikh Mohammed bin Zayed Al Nahyan, Crown Prince of Abu Dhabi and Chairman of the IFHC and H.E. Mohammed Al Bowardi, Deputy Chairman of IFHC, for their support. This study was conducted under the guidance of Reneco International Wildlife Consultants LLC., a consulting company managing ECWP. We are thankful to Dr. F. Lacroix, managing director, and G. Leveque, project director, for their supervision. We sincerely thank all ECWP staff from the Ecology division who participated in data collection. We are thankful to Pascale Reding and Grégoire Liénart for producing our study area map. We are grateful to Doug Armstrong and two anonymous reviewers for their helpful and constructive comments on the manuscript, as well as to Dr Thomas Martin for improving the English text.

References

- Amundsen, T., Stokland, J.N., 1990. Egg size and parental quality influence nestling growth in the shag. *The Auk* 107, 410–413.
- Araki, H., Cooper, B., Blouin, M.S., 2007. Genetic Effects of Captive Breeding Cause a Rapid, Cumulative Fitness Decline in the Wild. *Science* 318, 100–103.

502 Armstrong, D.P., Ewen, J.G., 2001. Assessing the value of follow-up translocations: a case
 503 study using New Zealand robins. *Biological Conservation* 101, 239–247.
 504 doi:10.1016/S0006-3207(01)00071-4
 505 Armstrong, D.P., Seddon, P.J., 2008. Directions in reintroduction biology. *Trends in Ecology*
 506 & *Evolution* 23, 20–25. <https://doi.org/10.1016/j.tree.2007.10.003>
 507 Armstrong, D.P., Le Coeur, C., Thorne, J.M., Panfylova, J., Lovegrove, T.G., Frost, P.G.H.,
 508 Ewen, J.G., 2017. Using Bayesian mark-recapture modelling to quantify the strength and
 509 duration of post-release effects in reintroduced populations. *Biological Conservation* 215,
 510 39–45. <https://doi.org/10.1016/j.biocon.2017.08.033>
 511 Azar, J.F., Rautureau, P., Lawrence, M., Calabuig, G., Hingrat, Y., 2016. Survival of
 512 reintroduced Asian houbara in United Arab Emirates' reserves. *The Journal of Wildlife*
 513 *Management* 80, 1031–1039. doi:10.1002/jwmg.21085
 514 Bacon, L., Hingrat, Y., Robert, A., 2017a. Evidence of reproductive senescence of released
 515 individuals in a reinforced bird population. *Biological Conservation* 215, 288–295.
 516 doi:10.1016/j.biocon.2017.08.023
 517 Bacon, L., Hingrat, Y., Jiguet, F., Monnet, A.C., Sarrazin, F., Robert, A., 2017b. Habitat
 518 suitability and demography, a time dependent relationship. *Ecol. Evol.* 2214–2222.
 519 Bacon, L., 2017. Etude des paramètres de reproduction et de la dynamique d'une population
 520 renforcée d'outardes Houbara nord-africaines (*Chlamydotis undulata undulata*) au
 521 Maroc. Thèse, Muséum National d'Histoire Naturelle, 192 pp.
 522 Barbanera, F., Pergams, O.R.W., Guerrini, M., Forcina, G., Panayides, P., Dini, F., 2010.
 523 Genetic consequences of intensive management in game birds. *Biological Conservation*
 524 143, 1259–1268. doi:10.1016/j.biocon.2010.02.035

525 Barron, D.G., Brawn, J.D., Weatherhead, P.J., 2010. Meta-analysis of transmitter effects on
 526 avian behaviour and ecology. *Methods in Ecology and Evolution* 1, 180–187.
 527 doi:10.1111/j.2041-210X.2010.00013.x
 528 Barton, K., 2015. MuMIn: multi-model inference. R package version 1 (9), 13.
 529 Bates, D., M. Martin, B. Bolker, S. Walker., 2015. Fitting Linear Mixed-Effects Models Using
 530 lme4. *Journal of Statistical Software* 67, 1–48 doi:10.18637/jss.v067.i01.
 531 Bertolero, A., Oro, D., Besnard, A., 2007. Assessing the efficacy of reintroduction programmes
 532 by modelling adult survival: the example of Hermann’s tortoise. *Animal Conservation*
 533 10, 360–368. doi:10.1111/j.1469-1795.2007.00121.x
 534 Bertolero, A., Oro, D., 2009. Conservation diagnosis of reintroducing Mediterranean pond
 535 turtles: what is wrong? *Animal Conservation* 12, 581–591. doi:10.1111/j.1469-
 536 1795.2009.00284.x
 537 Blomqvist, D., Johansson, O.C., Götmark, F., 1997. Parental quality and egg size affect chick
 538 survival in a precocial bird, the lapwing *Vanellus vanellus*. *Oecologia* 110, 18–24.
 539 doi:10.1007/s004420050128
 540 Bourass, K., Hingrat, Y., 2015. Diet of released captive-bred North-African houbara bustards.
 541 *European Journal of Wildlife Research* 61, 563–574. doi:10.1007/s10344-015-0930-8
 542 Bourass, K., Zaime, A., Qninba, A., Benhoussa, A., Rguibi, H., Hingrat, Y., 2012. Evolution
 543 saisonnière du régime alimentaire de l’Outarde houbara nord-africain, *Chlamydotis*
 544 *undulata*. *Bulletin de l’Institut Scientifique, Rabat* 34, 29–43.
 545 Brown, J.L., Collopy, M.W., Gott, E.J., Juergens, P.W., Montoya, A.B., Hunt, W.G., 2006.
 546 Wild-reared aplomado falcons survive and recruit at higher rates than hacked falcons in
 547 a common environment. *Biological Conservation* 131, 453–458.
 548 doi:10.1016/j.biocon.2006.02.021

549 Buner, F., Schaub, M., 2008. How do different releasing techniques affect the survival of
 550 reintroduced grey partridges *Perdix perdix*? *Wildlife Biology* 14, 26–35.
 551 doi:10.2981/0909-6396(2008)14[26:HDDRTA]2.0.CO;2

552 Buner, F.D., Browne, S.J., Aebischer, N.J., 2011. Experimental assessment of release methods
 553 for the re-establishment of a red-listed galliform, the grey partridge (*Perdix perdix*).
 554 *Biological Conservation* 144, 593–601. doi:10.1016/j.biocon.2010.10.017

555 Burnham, K.P., Anderson, D.E., 2002. Model selection and multimodel inference - A practical
 556 information-theoretic approach, Second Edition. Ed. Springer.

557 Cam, E., Aubry, L.M., Authier, M., 2016. The Conundrum of Heterogeneities in Life History
 558 Studies. *Trends in Ecology & Evolution* 31, 872–886. doi:10.1016/j.tree.2016.08.002

559 Casas, F., Benítez-López, A., García, J.T., Martín, C.A., Viñuela, J., Mougeot, F., 2015.
 560 Assessing the short-term effects of capture, handling and tagging of sandgrouse. *Ibis* 157,
 561 115–124. doi:10.1111/ibi.12222

562 Champagnon, J., Guillemain, M., Elmberg, J., Massez, G., Cavallo, F., Gauthier-Clerc, M.,
 563 2012. Low survival after release into the wild: assessing “the burden of captivity” on
 564 Mallard physiology and behaviour. *European Journal of Wildlife Research* 58, 255–267.
 565 doi:10.1007/s10344-011-0573-3

566 Chargé, R., Saint Jalme, M., Lacroix, F., Cadet, A., Sorci, G., 2010. Male health status,
 567 signalled by courtship display, reveals ejaculate quality and hatching success in a lekking
 568 species. *Journal of Animal Ecology* 79, 843–850. doi:10.1111/j.1365-2656.2010.01696.x

569 Chargé, R., Sorci, G., Saint Jalme, M., Lesobre, L., Hingrat, Y., Lacroix, F., Teplitsky, C., 2014.
 570 Does recognized genetic management in supportive breeding prevent genetic changes in
 571 life-history traits? *Evolutionary Applications* 7, 521–532. doi:10.1111/eva.12150

572 Christensen, R. H. B., 2015. ordinal - Regression Models for Ordinal Data. R package version
 573 2015, 6–28

574 Converse, S.J., Moore, C.T., Armstrong, D.P., 2013. Demographics of reintroduced
 575 populations: Estimation, modelling, and decision analysis. *Journal of Wildlife*
 576 *Management* 77, 1081–1093. <https://doi.org/10.1002/jwmg.590>
 577 Curio, E., 1983. Why young birds reproduce less well? *Ibis* 125, 400–404. doi:10.1111/j.1474-
 578 919X.1983.tb03130.x
 579 Dickens, M.J., Delehanty, D.J., Michael Romero, L., 2010. Stress: An inevitable component of
 580 animal translocation. *Biological Conservation* 143, 1329–1341.
 581 doi:10.1016/j.biocon.2010.02.032
 582 Ewen, J.G., Armstrong D.P., Parker K.A., Seddon, P.J., 2012. *Reintroduction Biology:*
 583 *Integrating Science and Management*. Wiley-Blackwell
 584 Fischer, J., Lindenmayer, D.B., 2000. An assessment of the published results of animal
 585 relocations. *Biological Conservation* 96, 1–11. doi:10.1016/S0006-3207(00)00048-3
 586 Frankham, R., 2008. Genetic adaptation to captivity in species conservation programs.
 587 *Molecular Ecology* 17, 325–333. doi:10.1111/j.1365-294X.2007.03399.x
 588 Galbraith, H., 1988. Effects of egg size and composition on the size, quality and survival of
 589 lapwing *Vanellus vanellus* chicks. *Journal of Zoology* 214, 383–398. doi:10.1111/j.1469-
 590 7998.1988.tb03747.x
 591 Gaucher, P., 1995. Breeding biology of the houbara bustard *Chlamydotis undulata* in Algeria.
 592 *Alauda* 63, 291–298.
 593 Gelman, A., 2008. Scaling regression inputs by dividing by two standard deviations. *Statistics*
 594 *in medicine* 27, 2865–2873.
 595 Goriup, P. D., 1997. The world status of the houbara bustard *Chlamydotis undulata*. *Bird*
 596 *Conservation International* 7, 373–397.
 597 Grant, T.A., Shaffer, T.L., 2012. Time-specific patterns of nest survival for ducks and
 598 passerines breeding in North Dakota. *The Auk* 129, 1–10.

599 Hardouin, L.A., Nevoux, M., Robert, A., Gimenez, O., Lacroix, F., Hingrat, Y., 2012.
600 Determinants and costs of natal dispersal in a lekking species. *Oikos* 121, 804–812.
601 doi:10.1111/j.1600-0706.2012.20313.x

602 Hardouin, L.A., Robert, A., Nevoux, M., Gimenez, O., Lacroix, F., Hingrat, Y., 2014.
603 Meteorological conditions influence short-term survival and dispersal in a reinforced bird
604 population. *Journal Applied Ecology* 51, 1494–1503. doi:10.1111/1365-2664.12302

605 Hardouin, L.A., Hingrat, Y., Nevoux, M., Lacroix, F., Robert, A., 2015a. Survival and
606 movement of translocated houbara bustards in a mixed conservation area. *Animal*
607 *conservation*, 18(5), 461–470.

608 Hardouin, L.A., Legagneux, P., Hingrat, Y., Robert, A., 2015b. Sex-specific dispersal responses
609 to inbreeding and kinship. *Animal Behaviour* 105, 1–10.
610 doi:10.1016/j.anbehav.2015.04.002

611 Herzog, M., 2009. nest survival. R package that performs logistic exposure nest survival
612 analyses.

613 Hill, D., Robertson, P., 1988. Breeding Success of Wild and Hand-Reared Ring-Necked
614 Pheasants. *The Journal of Wildlife Management* 52, 446–450. doi:10.2307/3801588

615 Hoyt, D., 1979. Practical methods of estimating volume and fresh weight of birg eggs. *The Auk*
616 96, 73–77.

617 Huff, D.D., Miller, L.M., Chizinski, C.J., Vondracek, B., 2011. Mixed-source reintroductions
618 lead to outbreeding depression in second-generation descendents of a native North
619 American fish. *Molecular Ecology* 20, 4246–4258. doi:10.1111/j.1365-
620 294X.2011.05271.x

621 IUCN/SSC. 2013. Guidelines for reintroductions and other conservation translocations. *Version*
622 *1.0. Gland, Switzerland. Available from*
623 http://www.issg.org/pdf/publications/RSG_ISSG-Reintroduction-Guidelines-2013.pdf

624 Koshkin, M., Burnside, R.J., Packman, C.E., Collar, N.J., Dolman, P.M., 2016. Effects of
625 habitat and livestock on nest productivity of the Asian houbara *Chlamydotis macqueenii*
626 in Bukhara Province, Uzbekistan. *European Journal Wildlife Research* 62, 447–459.
627 doi:10.1007/s10344-016-1018-9

628 Lacroix, F., Seabury, J., Al Bowardi, M., and Renaud, J., 2003. The Emirates Center for
629 Wildlife Propagation: comprehensive strategy to secure self-sustaining wild populations
630 of houbara bustard (*Chlamydotis undulata undulata*) in Eastern Morocco. *Houbara News*
631 5, 60–62.

632 Le Gouar, P., Robert, A., Choisy, J.-P., Henriquet, S., Lecuyer, P., Tessier, C., Sarrazin, F.,
633 2008. Roles of Survival and Dispersal in Reintroduction Success of Griffon Vulture
634 (*Gyps fulvus*). *Ecological Applications* 18, 859–872. doi:10.1890/07-0854.1

635 Le Maho, Y., Whittington, J.D., Hanuise, N., Pereira, L., Boureau, M., Brucker, M., Chatelain,
636 N., Courtecuisse, J., Crenner, F., Friess, B., Grosbellet, E., Kernaléguen, L., Olivier, F.,
637 Saraux, C., Vetter, N., Viblanc, V.A., Thierry, B., Tremblay, P., Groscolas, R., Le Bohec,
638 C., 2014. Rovers minimize human disturbance in research on wild animals. *Nature*
639 *Methods* 11, 1242–1244. <https://doi.org/10.1038/nmeth.3173>

640 Lesobre, L., Lacroix, F., Caizergues, A., Hingrat, Y., Chalah, T., Jalme, M.S., 2010.
641 Conservation genetics of Houbara Bustard (*Chlamydotis undulata undulata*): population
642 structure and its implications for the reinforcement of wild populations. *Conservation*
643 *Genetics* 11, 1489–1497. doi:10.1007/s10592-009-9979-9

644 Mabee, T.J., 1997. Using Eggshell Evidence to Determine Nest Fate of Shorebirds. *The Wilson*
645 *Bulletin* 109, 307–313. doi:10.1111/j.1557-9263.2006.00037.x

646 Manolis, J.C., Andersen, D.E., Cuthbert, F.J., 2000. Uncertain nest fates in songbird studies and
647 variation in mayfield estimation. *The Auk* 117, 615–626. doi:10.1642/0004-
648 8038(2000)117[0615:UNFISS]2.0.CO;2

649 Martin, T.E., 1995. Avian Life History Evolution in Relation to Nest Sites, Nest Predation, and
 650 Food. Ecological Monographs 65, 101–127. doi:10.2307/2937160
 651 Mauck, R.A., Huntington, C.E., Doherty Jr, P.F., 2012. Experience versus effort: what explains
 652 dynamic heterogeneity with respect to age? Oikos 121, 1379–1390. doi:10.1111/j.1600-
 653 0706.2012.20271.x
 654 Mayfield, H.F., 1975. Suggestions for Calculating Nest Success. The Wilson Bulletin 87, 456–
 655 466. doi:10.2307/4160682
 656 Mihoub, J.-B., Robert, A., Gouar, P.L., Sarrazin, F., 2011. Post-Release Dispersal in Animal
 657 Translocations: Social Attraction and the “Vacuum Effect”. PLOS ONE 6, e27453.
 658 doi:10.1371/journal.pone.0027453
 659 Monnet, A.-C., Hingrat, Y., Jiguet, F., 2015. The realized niche of captive-hatched Houbara
 660 Bustards translocated in Morocco meets expectations from the wild. Biological
 661 Conservation 186, 241–250. doi:10.1016/j.biocon.2015.03.013
 662 Montalvo, A.M., Ellstrand, N.C., 2001. Nonlocal transplantation and outbreeding depression in
 663 the subshrub *Lotus scoparius* (Fabaceae). American Journal of Botany 88, 258–269.
 664 Morales, M.B., Alonso, J.C., Alonso, J., 2002. Annual productivity and individual female
 665 reproductive success in a Great Bustard *Otis tarda* population. Ibis 144, 293–300.
 666 doi:10.1046/j.1474-919X.2002.00042.x
 667 Nussey, D.H., Kruuk, L.E.B., Donald, A., Fowlie, M., Clutton-Brock, T.H., 2006. The rate of
 668 senescence in maternal performance increases with early-life fecundity in red deer.
 669 Ecology Letters 9, 1342–1350. doi:10.1111/j.1461-0248.2006.00989.x
 670 Pérez, J.A., Sánchez-García, C., Díez, C., Bartolomé, D.J., Alonso, M.E., Gaudioso, V.R.,
 671 2015. Are parent-reared red-legged partridges (*Alectoris rufa*) better candidates for re-
 672 establishment purposes? Poultry Science 94, 2330–2338. doi:10.3382/ps/pev210

673 Ponjoan, A., Bota, G., De La Morena, E.L.G., Morales, M.B., Wolff, A., Marco, I., Mañosa, S.,
 674 2008. Adverse Effects of Capture and Handling Little Bustard. *The Journal of Wildlife*
 675 *Management* 72, 315–319. doi:10.2193/2006-443
 676 R Development Core Team, 2017. R: A language and environment for statistical computing. R
 677 Foundation for Statistical Computing, Vienna, Austria.
 678 Robert, A., Couvet, D., Sarrazin, F., 2007. Integration of demography and genetics in
 679 population restorations. *Ecoscience* 14, 463–471. doi:10.2980/1195-
 680 6860(2007)14[463:IODAGI]2.0.CO;2
 681 Robert, A., 2009. Captive breeding genetics and reintroduction success. *Biological*
 682 *Conservation* 142, 2915–2922. doi:10.1016/j.biocon.2009.07.016
 683 Robert, A., Colas, B., Guigon, I., Kerbiriou, C., Mihoub, J.-B., Saint-Jalme, M., Sarrazin, F.,
 684 2015a. Defining reintroduction success using IUCN criteria for threatened species: a
 685 demographic assessment. *Animal Conservation* 18, 397–406. doi:10.1111/acv.12188
 686 Robert, A., Bolton, M., Jiguet, F., Bried, J., 2015b. The survival–reproduction association
 687 becomes stronger when conditions are good. *Proceedings of the Royal Society B* 282,
 688 20151529. doi:10.1098/rspb.2015.1529
 689 Roche, E.A., Cuthbert, F.J., Arnold, T.W., 2008. Relative fitness of wild and captive-reared
 690 piping plovers: Does egg salvage contribute to recovery of the endangered Great Lakes
 691 population? *Biological Conservation* 141, 3079–3088. doi:10.1016/j.biocon.2008.09.014
 692 Saint Jalme, M., Van Heezik, Y., 1996. Propagation of the houbara bustard. Paul Keegan,
 693 London.
 694 Sánchez-García, C., Alonso, M.E., Pérez, J.A., Rodríguez, P.L., Gaudioso, V.R., 2011.
 695 Comparing fostering success between wild-caught and game farm bred captive red-
 696 legged partridges (*Alectoris rufa*, L.). *Applied Animal Behaviour Science* 133, 70–77.
 697 doi:10.1016/j.applanim.2011.04.012

698 Sarrazin, F., Bagnolini, C., Pinna, J.L., Danchin, E., Clobert, J., 1994. High Survival Estimates
699 of Griffon Vultures (*Gyps fulvus fulvus*) in a Reintroduced Population. *The Auk* 111,
700 853–862. doi:10.2307/4088817

701 Sarrazin, F., Bagnolin, C., Pinna, J.L., Danchin, E., 1996. Breeding biology during
702 establishment of a reintroduced Griffon Vulture *Gyps fulvus* population. *Ibis* 138, 315–
703 325. doi:10.1111/j.1474-919X.1996.tb04344.x

704 Shaffer, T.L., 2004. A unified approach to analyzing nest success. *The Auk* 121, 526–540.

705 Sutherland, W.J., Armstrong, D., Butchart, S.H.M., Earnhardt, J.M., Ewen, J., Jamieson, I.,
706 Jones, C.G., Lee, R., Newbery, P., Nichols, J.D., Parker, K.A., Sarrazin, F., Seddon, P.J.,
707 Shah, N., Tatayah, V., 2010. Standards for documenting and monitoring bird
708 reintroduction projects. *Conservation Letters* 3, 229–235. doi:10.1111/j.1755-
709 263X.2010.00113.x

710 Tavecchia, G., Viedma, C., Martínez-Abraín, A., Bartolomé, M.-A., Gómez, J.A., Oro, D.,
711 2009. Maximizing re-introduction success: Assessing the immediate cost of release in a
712 threatened waterfowl. *Biological Conservation* 142, 3005–3012.
713 doi:10.1016/j.biocon.2009.07.035

714 van de Pol, M., Verhulst, S., 2006. Age- Dependent Traits: A New Statistical Model to Separate
715 Within- and Between- Individual Effects. *The American Naturalist* 167, 766–773.
716 doi:10.1086/503331

717 Williams, T.D., 1994. Intraspecific variation in egg size and egg composition in birds: effects
718 on offspring fitness. *Biological Reviews of the Cambridge Philosophical Society* 69, 35–
719 59.

720 Zuur, A.F., Ieno, E.N., Walker, N., Saveliev, A.A., Smith, G.M., 2009. Mixed effects models
721 and extensions in ecology with R, Statistics for Biology and Health. Springer New York,
722 New York, NY. doi:10.1007/978-0-387-87458-6

723

724